

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|-----------|--|
| (51) International Patent Classification ⁶ : C12N 9/24, A23L 1/185, 1/10, C07K 14/415 | A1 | (11) International Publication Number: WO 98/49278 (43) International Publication Date: 5 November 1998 (05.11.98) |
| (21) International Application Number: PCT/EP98/02590 (22) International Filing Date: 4 May 1998 (04.05.98) (30) Priority Data: 97870060.7 30 April 1997 (30.04.97) EP (34) Countries for which the regional or international application was filed: BE et al. (71) Applicant (for all designated States except US): K.U. LEUVEN RESEARCH & DEVELOPMENT [BE/BE]; Groot Begijnhof 59, B-3000 Leuven (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): DEBYSER, Winok [BE/BE]; Willem Coosemansstraat 23, B-3010 Kessel-Lo (BE). DELCOUR, Jan [BE/BE]; Kastanjelaan 14, B-3001 Leuven (BE). (74) Agents: VAN MALDEREN, Joëlle et al.; Office van Malderen, Place Reine Fabiola 6/1, B-1083 Brussels (BE). | | (81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: INHIBITORS OF CELLULOLYTIC, XYLANOLYTIC AND β -GLUCANOLYTIC ENZYMES (57) Abstract The present invention concerns an inhibitor of xylanolytic and/or β -glucanolytic enzymes, method for obtaining the inhibitor, said inhibitor and processes for obtaining micro-organism, plant or plant material wherein the activity of the inhibitor according to the invention is increased or reduced and to the use of the inhibitor, the cited micro-organism, plant or plant material in a variety of processes and applications. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

5

INHIBITORS OF CELLULOLYTIC, XYLANOLYTIC AND β -GLUCANOLYTIC
ENZYMES

10 Field of the invention

This invention relates to an inhibitor of cellulolytic, xylanolytic and/or β -glucanolytic enzymes (sometimes also referred to as cellulases, pentosanases and/or hemicellulases) especially an inhibitor of pentosan
15 degrading enzymes such as endoxylanase (such as EC: 3.2.1.8), β -xylosidase (such as EC: 3.2.1.37), and α -L-arabinofuranosidase (such as EC: 3.2.1.55), to inhibitors of cellulase (such as EC: 3.2.1.4), β -glucanase (such as EC: 3.2.3.73 or EC: 3.2.1.6), and to inhibitors of other
20 xylan, arabinoxylan and β -glucan degrading enzymes, which are present in micro-organisms, plants, plant materials or fractions thereof, (such as cereals, cereal grains, cereal flours or fractions thereof).

The present invention is also related to a
25 method for obtaining said inhibitor, as well as to the use of said inhibitor in different areas of food, feed and/or beverage technologies, such as malting and brewing, the production of animal feedstuffs such as to increase their conversion, the production of baked and/or extruded cereal
30 products such as straight dough, sponge and dough and Chorleywood breads, breakfast cereals, different types of

biscuits, pasta and noodles, the production of starch derived syrups, sorbitol, xylose and/or xylitol, the wheat gluten-starch separation industry, maize processing, the improvement of plant disease resistance, in nutraceutical
5 or pharmaceutical applications such as maintaining the structure of dietary fiber material, and in the field of paper and pulp technologies.

Background of the invention

10 Apart from barley malt, unmalted cereals such as wheat are commonly used in beer production (Pierce, J.S., *Proceedings of the European Brewery Convention Congress, Madrid, 1987*, 445). Unmalted wheat (40-50%) is e.g. used for the production of Belgian white (wheat)
15 beers.

 Although barley and wheat endosperm cell walls contain 20 and 70% (w/w) arabinoxylan respectively (Ballance, G.M., & Manners, D.J., *Carbohydrate Research*, 1978, 61,107; Fincher, G.B., & Stone, B.A. In: *Advances in*
20 *Cereal Science and Technology*, Vol. VIII. Y. Pomeranz, (Ed), Am. Assoc. Cereal Chem., St. Paul (MN), 1986, 207), their total arabinoxylan content is comparable, i.e. 2.8 to 7.1% (w/w) for barley and 3.6 to 7.1% (w/w) for wheat (Henry, J., *Journal of the Science of Food and Agriculture*,
25 1985, 36, 1243; Hashimoto, S., Shogren, M.D. & Pomeranz, Y., *Cereal Chemistry*, 1987, 64, 30).

 The grains also contain comparable levels of water-extractable arabinoxylan, i.e. 0.24 to 0.80% (w/w) for barley and 0.25 to 1.18% for wheat (Henry, J., *Journal*
30 *of the Science of Food and Agriculture*, 1985, 36, 1243; Hashimoto, S., Shogren, M.D. & Pomeranz, Y., *Cereal Chemistry*, 1987, 64, 30; Åman, P., & Hesselman, K., *Swedish*

Journal of Agricultural Research, 1984, 14, 135; Girhammer, U., & Nair, B.M., *Food Hydrocolloids*, 1992, 6, 285). Furthermore, barley and wheat endosperm cell walls contain 70 and 20% β -glucan respectively (Ballance, G.M., & 5 Manners, D.J., *Carbohydrate Research*, 1978, 61, 107; Fincher, G.B., & Stone, B.A. In: *Advances in Cereal Sciences and Technology*, Vol. VIII. Y. Pomeranz, (Ed), Am. Assoc. Cereal Chem., St. Paul (MN), 1986, 207).

Barley grains contain 1.7 to 4.1% (w/w) 10 water-extractable and 3.6 to 6.4% (w/w) total β -glucan (Anderson, M.A., Cook, J.A., & Stone, B.A., *Journal of the Institute of Brewing*, 1978, 84, 233-239; Henry, J., *Journal of the Science of Food and Agriculture*, 1985, 36, 1243). Wheat grains contain 0.1 to 0.8% (w/w) water-extractable 15 and 0.6 to 1.4% (w/w) total β -glucan (Anderson, M.A., Cook, J.A., & Stone, B.A., *Journal of the Institute of Brewing*, 1978, 84, 233-239; Henry, J., *Journal of the Science of Food and Agriculture*, 1985, 36, 1243). As in wheat only low levels of arabinoxylan (Cleemput, G., Bleukx, W., van Oort, 20 M., Hessing, M. & Delcour, J.A., *Journal of Cereal Science*, 1995, 22, 139) and β -glucan degrading enzyme activities can be measured, the barley malt must be mostly responsible for wheat and malt arabinoxylan and β -glucan hydrolysis during brewing.

25 Efficient hydrolysis of arabinoxylans and β -glucan is important because such compounds can be involved in production problems such as wort viscosity (Ducroo, P. & Frelon, P.G., *Proceedings of the European Brewery Convention Congress*, Zurich, 1989, 445; Viëtor, R.J. & 30 Voragen, A.G.J., *Journal of the Institute of Brewing*, 1993, 99, 243) and filterability and haze formation (Coote, N. &

Kirsop, B.H. 1976., *Journal of the Institute of Brewing*, 1976, 82, 34; Izawa, M., Kano, Y. & Kanimura, M. 1991. *Proceedings Aviemore Conference on Malting, Brewing and Distillling*, 1990, 427).

5 In other areas efficient hydrolysis of xylans and/or arabinoxylans is highly desirable as well. Examples include rye and wheat breadmaking processes, paper and pulp technologies. It follows that a lot of research efforts have been devoted to the (potential) applications of xylan
10 and/or arabinoxylan hydrolysis enzymes due to their applications as described above.

Summary of the invention

 The present invention concerns an inhibitor
15 of cellulolytic, xylanolytic and/or β -glucanolytic enzymes, preferably an inhibitor of endoxylanase, of β -glucanase, of β -xylosidase, of α -L-arabinofuranosidase, and of other xylan, arabinoxylan and β -glucan degrading enzymes preferably obtained from micro-organisms, plants, plant
20 materials or fractions thereof (such as cereals, cereal grains, cereal germs or fractions thereof, cereal flours or fractions thereof).

 "An inhibitor of an enzyme" means a molecule which is able to inhibit partially or totally the activity
25 of said enzyme. In irreversible inhibition, the inhibitor is covalently linked to the enzyme or bound so tightly that its dissociation from the enzyme is very slow. In this case, the inhibitor usually mimicks the normal substrate of said enzyme in a cross-linking reaction. In contrast,
30 reversible inhibition may be characterised by a rapid equilibrium between the enzyme and the inhibitor. A

competitive inhibitor prevents the substrate from binding to the active site and may reduce the reaction rate by diminishing the proportion of enzyme molecules that are bound to substrate. In non-competitive inhibition, the inhibitor may decrease the turnover number. Competitive inhibition can be distinguished from non-competitive inhibition by determining whether the inhibition can be overcome by raising the substrate concentration. Inhibitors isolated from a specific biological species and that are of proteinaceous or glycoproteinaceous nature can be active against enzymes of the same species (i.e. endogeneous enzymes) and/or against enzymes of different species (i.e. exogeneous enzymes).

Advantageously the inhibitor of the invention can be produced by micro-organisms or may be present in various extraction media from micro-organisms or plant material, such as cereals or fractions thereof, such as cereal grains or fractions thereof, such as cereal germs or fractions thereof, such as cereal flours or fractions thereof, such as from wheat, durum wheat, rye, triticale, barley, sorghum, oats, maize and/or rice, from which it can be obtained by the methods well known by the person skilled in the art. According to a preferred embodiment of the present invention, the inhibitor is a xylanase inhibitor which is typically water-soluble alkaline proteinaceous species, having a pI (i.e. -log of the isoelectric point) of greater than about 7.0. The xylanase inhibitor molecular weight as determined by SDS-page is typically 40-43 kDa. Following reduction with β -mercaptoethanol three SDS-page protein bands are found with SDS-page molecular weights of ca. 40-43 kDa, ca. 30 kDa, and ca. 10 kDa. The N-terminal sequence of the 40-43 kDa protein or glycoprotein has not

been described until now and is typically as follows: SEQ ID No. 1: Lys-Gly-Leu-Pro-Val-Leu-Ala-Pro-Val-Thr-Lys-Xaa-Thr-Ala, wherein Xaa being preferably Asp. The 30 kDa band has the above described typical N-terminal amino acid SEQ ID No.1, while the N-terminal amino acid sequence of the 10 kDa band is typically as follows: SEQ ID No. 2: Xaa-Ala-Pro-Val-Ala-Lys-Met-Val-Leu-Pro-Val-Ala-Met-Lys-Glu-Xaa-Val, wherein the first Xaa being preferably Ser, Phe, or Gly, and wherein the second Xaa is unidentified. This sequence has not been described before.

Therefore, the present invention is also related to an inhibitor with an SDS-page molecular weight of typically 40-43 kDa being a protein or glycoprotein having a marker whose amino acid sequence has more than 70% homology, preferably more than 85% homology, more preferably is identical with SEQ ID No. 1.

The present invention is furthermore also related to an inhibitor with an SDS-page molecular weight of typically 30 kDa being a protein or glycoprotein having a marker whose amino acid sequence has more than 70% homology, preferably more than 85% homology, more preferably is identical with SEQ ID No. 1.

The present invention is furthermore also related to an inhibitor with an SDS-page molecular weight of typically 10 kDa being a protein or glycoprotein having a marker whose amino acid sequence has more than 70% homology, preferably more than 85% homology, more preferably is identical with SEQ ID No. 2.

Advantageously, said markers are the end-terminal amino acid sequences of the protein or glycoprotein.

According to the invention, a marker of a

protein or glycoprotein means a specific amino acid sequence (or its corresponding nucleotide acid sequence) that is able to distinguish one protein family from another protein family.

5 The inhibitory effect towards xylan and/or arabinoxylan hydrolysing enzymes can be e.g. demonstrated by the endoxylanase method with AZCL arabinoxylan (cfr. infra). Likewise, the inhibitory effect towards β -glucan hydrolysing enzymes can be e.g. demonstrated by the β -
10 glucanase method with AZCL β -glucan (cfr. infra).

 The invention also relates to a method for obtaining said inhibitor from a micro-organism, such as a genetically modified micro-organism which expresses said inhibitors, from a plant, or from a plant material such as
15 cereals, cereal grains, cereal flours or fractions thereof), by subjecting said plant, said plant material and/or said micro-organism to one or more extraction and/or fractionation steps.

 Another aspect of the present invention is
20 related to a method for genetically transforming a micro-organism, a plant or a plant material in order to obtain the expression of the inhibitor according to the invention wherein the micro-organism, the plant or plant material is genetically modified by the introduction of a genetic
25 material encoding said inhibitor into the micro-organism, the plant or plant material and obtain its translation and expression by genetic engineering methods well known by the man skilled in the art.

 The invention furthermore relates to
30 processes aiming at changing, preferably reducing or increasing level of said inhibitor in a micro-organism, a plant or a plant material, by reducing or increasing the

expression of said inhibitor, by the methods well known by the man skilled in the art and/or by using molecules which are able to block the inhibitor activity or activate said inhibitor.

5 The invention furthermore relates to the obtained inhibitor, micro-organism, plant, plant material, and/or fractions thereof and to their use in different areas of food, feed and/or beverage technologies, such as improving malting and brewing, improving animal feedstuffs
10 efficiency, baked and/or extruded cereal products (such as straight dough, sponge and dough and Chorleywood breads, breakfast cereals, different types of biscuits, pasta and noodles), improving the production of starch derived syrups, sorbitol, xylose and/or xylitol, improving wheat
15 gluten-starch separation and production, maize processing, improving plant disease resistance, improving nutraceutical or pharmaceutical applications (such as maintaining the structure of dietary fiber material), and improving paper and pulp technologies.

20 The present invention will be described in details in the following description of a preferred embodiment without limiting the scope of the present invention.

25 Detailed description of the invention

 During the course of their work dealing with the structure of arabinoxylans in Belgian white beers and in intermediates in the production process, the inventors unexpectedly found indications for inhibition of the
30 xylanolytic barley malt system by wheat water extractables. This has not been reported before, although it has clearly been established that endogenous and exogenous α -amylase

(Deponte, R., Parlamenti, T., Petrucci, V., Silano, V., & Tomasi, M., *Cereal Chemistry*, 1976, 53, 805; Buonocore, V., Petrucci, T., & Silano, V., *Phytochemistry*, 1977, 16, 811; Mundy, J., Hejgaard, J., & Svendsen, I., *Federation of Societies*, 1984, 167, 210; Silano, V. α -Amylase inhibitors. In: *Enzymes and their Role in Cereal Technology*, J.E. Kruger, D. Lineback and C.E. Stauffer, (Eds). Am. Assoc. Cereal Chem., St. Paul (MN), 1987, 141) and protease (Birk, Y., *Methods Enzymology*, 1976, 45, 723; Lawszkowski, M., & Kato, I., *Annual Review of Biochemistry*, 1980, 49, 593) inhibitors are present in cereal grains.

Indeed, when one measured the solubilization of arabinoxylans during brewing with barley malt and unmalted wheat, with the objectives (1) to relate enzymic activities of the starting materials with the arabinoxylan contents of corresponding worts, and (2) to investigate in which way wheat interferes with the solubilization of arabinoxylans during brewing, there was evidence for the presence of xylanase inhibitors in wheat. This was indeed observed when one compared the solubilization of arabinoxylans and the release of free xylose (Xyl) in wort prepared with 60% malt and 40% wheat with that in a 100% malt wort.

Under certain experimental conditions, the addition to the wort of a xylanase of microbial origin clearly improved arabinoxylan solubilization during wort preparation.

Examples

30 Materials

β -D-Allose, β -mercaptoethanol, p-nitro-phenyl- β -D-xylopyranoside and Trizma base (reagent grade,

tris[hydroxymethyl]amino-methane) were obtained from Sigma, St-Louis, MO, USA. Azurine-crosslinked (AZCL) wheat arabinoxylan (Xylazyme arabinoxylan tablets), AZCL and Xylanase M4 from *Aspergillus niger* was from Megazyme, Bray, Ireland. Microbial xylanases from the micro-organisms *Bacillus subtilis*, *Trichoderma viride* and *Aspergillus niger* were obtained from NV Puratos, Groot-Bijgaarden, Belgium. Buffer A was: 0.025 M sodium acetate, pH 4.7; Buffer B was: 0.025 M sodium maleate, pH 6.0; Buffer C was: 0.025 M sodium phosphate pH 6.0; Buffer D was: 0.250 mM sodium acetate pH 5.0; Buffer E was: 0.025 M sodium acetate pH 5.0.

Barley malt samples were supplied by Cargill Malt Division, Herent (Belgium). The inventors used a two-rowed winter barley variety (Clarine) with a low endoxylanase activity and low water-extractable Xyl content, and two malts from a six-rowed winter barley variety Plaisant, with a high water-extractable Xyl content. Plaisant malt samples 1 and 2 had high and low endoxylanase activities respectively. Wheat samples were from Amylum, Aalst (Belgium) and SAPSA SES SA, Jodoigne (Belgium). The inventors used Skirlou and Soissons with high and low water-extractable Xyl contents, respectively. Wheat germs were supplied by Ceres, Vilvoorde (Belgium). Rye flour was from a mixture of Dutch rye varieties supplied by Meneba, Rotterdam (The Netherlands). Barley from the variety Clarine was supplied by Cargill, Malt Division, Herent (Belgium). Clarine barley, Plaisant 1 and Plaisant 2 barley malts, Skirlou and Soissons wheat wholemeals were prepared either with the Tecator sample mill (Höganäs, Sweden) or for the brewing experiments with an EBC-approved laboratory mill (Analytica-EBC, Fourth

edition, Brauerei- und Getränke- Rundschau, Zurich, 1987). Soissons wheat flour was produced with a Bühler MLU-202 laboratory mill (Bühler, Uzwil, Switzerland, extraction yield 70%).

5

Extracts

BMWM1 WWM, and WG

Samples (3.00 g) of ground barley malt and wheat or wheat germ (1.00 g) were suspended in buffer A
10 (10.0 mL). After 15 min of vigorous shaking at room temperature, the suspensions were centrifuged (3,000 g, 15 min, 20°C). The resulting extracts are referred to as BMWM1 (barley malt wholemeal extract 1), WWM (wheat wholemeal extract), and WG (wheat germ extract).

15 *WF, RF, and BWM*

Samples of the appropriate flour or wholemeal (2.50 g) were suspended in buffer B (10.0 mL). After 15 min of vigorous shaking at room temperature, the suspensions were centrifuged (10,000 g, 15 min, 20°C) and the
20 supernatants were filtered (0.45 µ). The resulting wholemeal extracts are referred to as WF (wheat flour extract), RF (rye flour extract), and BWM (barley wholemeal extract).

BMWM2

25 Samples (5.00 g) of ground barley malt were suspended in buffer B (10.0 mL). After 15 min of vigorous shaking at room temperature, the suspensions were centrifuged (10,000 g, 15 min, 20°C) and the supernatants were filtered (0.45 µ). The resulting wholemeal extracts
30 are referred to as BMWM2 (barley malt wholemeal extract 2).

Methods

Determination of Xyl content

Extraction and hydrolysis procedures were as described by Cleemput et al (Cleemput, G., Roels, S.P., van Oort, M., Grobet P.J. & Delcour, J.A., *Cereal Chemistry*, 1993, 70, 324), with heating (130°C) of samples of whole meal (wheat and barley malt) for 5 hours to eliminate enzyme activity prior to extraction. Worts were analysed in the same way as the water-extracts of the whole meal flours. Free Xyl was determined by omitting the hydrolysis step prior to alditol acetate preparation. Alditol acetate samples (1µl) (Englyst, H.N. & Cummings J.H., *Analyst*, 1984, 109, 937) were separated at 225°C on a Supelco SP-2380 column (30 m, 0.32 mm ID, 0.2 µm film thickness) and detected with a flame ionisation detector in a Chrompack 9011 Chromatograph (Middelburg, The Netherlands). Injection and detection temperatures were 275°C. β-D-Allose was used as internal standard. The arabinose (Ara) measured originated from both arabinoxylan and arabinogalactan making it impossible to calculate arabinoxylan levels as $0.88 \times (\text{Ara} + \text{Xyl})$ (Cleemput, G., van Oort, M., Hessing, M., Bergmans, M.E.F., Gruppen, H., Grobet, P.J., Delcour, J.A., *Journal of Cereal Science*, 1995, 22, 73-84). Moreover, as in water-extracts of wheat wholemeal a substantial part of the galactose (Gal) does not stem from arabinogalactan, correction of Ara figures for arabinogalactan by assuming that the Gal/Ara ratio in arabinogalactan is 1.5 as known for wheat flours (Izydorczyk, M., Biliaderis, C.G. & Bushuk, W., *Cereal Chemistry*, 1991, 68, 139-144) was equally impossible. In what follows, therefore, the Xyl figures are used as a relative measure for arabinoxylans levels. In a similar

way, the increase in Xyl levels during brewing is indicative of arabinoxylan solubilization during brewing.

Measurement of endoxylanase (EC 3.2.1.8) activity and
5 inhibition thereof

Extracts (1.0 mL) BMWM1 and WWM (cfr. supra) were incubated for 5 min at 50°C, before adding an AZCL-xylan tablet (Megazyme). The incubation was then continued for 60 min at 50°C. The reaction was terminated by adding
10 1% (w/v) Trizma base (10.0 mL) and vigorous vortex stirring. After 5 min at room temperature, the tubes were shaken vigorously and the contents filtered through a Whatman N°1 filter. The absorbance was measured at 590 nm against a control, which was prepared by incubating the
15 extract without the substrate tablet for 60 min at 50°C. The substrate tablet was added after adding 1% (w/v) Trizma base to the extract. The activity was expressed as the difference in the absorbance at 590 nm between the sample and control and expressed per gram dry malt ($\Delta A_{590}/g$).

20 The endoxylanase activity of 0.6 mL BMWM1 (cfr. supra) and 0.4 mL buffer A was compared with the activity of 0.6 mL BMWM1 to which 0.4 mL WWM was added. In some cases, the WWM was boiled for 30 min and centrifuged (3,000 g, 15 min, 20°C) prior to addition.

25 In the evaluation of the inhibition of microbial enzymes by extracts from different cereals the following procedure was used. Extracts (WF, WG, RF, and BWM) or boiled (30 min, 100°C) and centrifuged (10,000 g, 15 min, 20°C) extracts (250 μ L) were preincubated for 30
30 min at room temperature with 250 μ L of appropriately diluted microbial xylanase solution, the xylazyme tablet was added and the mixture was incubated for 60 min at 50°C.

The remainder of the procedure was as described above with addition of 5.0 mL 2% (w/v) Trizma base instead of 10.0 mL 1% (w/v) to terminate the reaction.

5 *Measurement of β -glucanase (EC: 3.2.3.73) activity and inhibition thereof*

Extracts (WF, RF, and BWM) or boiled (30 min, 100°C) and centrifuged (10,000 g, 15 min, 20°C) extracts (450 μ L) were preincubated for 30 min at room temperature
10 with 50 μ L of BMW2, the β -glucanase tablet was added and the mixture was incubated for 60 min at 40°C. The remainder of the procedure was as described above with addition of 5.0 mL 2% (w/v) Trizma base instead of 10.0 mL 1% (w/v) to terminate the reaction.

15

Brewing

Worts were prepared in duplicate according to the EBC method (Analytica-EBC, Fourth edition, Brauerei- und Getränke- Rundschau, Zurich, 1987). For the 100% barley
20 malt worts, 50.0 g barley malt was used and for the worts with 40% wheat, 30.0 g barley malt and 20.0 g wheat. Worts were centrifuged for 15 min at 2,000 \times g (room temperature). The spent grains were washed (150 mL) and the washings were added to the worts.

25 The *Bacillus subtilis* endoxylanase was added to the water (46°C) before mixing with 60% Clarine malt and 40% Soissons or Skirlou wheat. The level of endoxylanase added to worts (0.867 $\Delta A_{590}/g$) was equal to that needed to increase the endoxylanase activity of Clarine malt (0.750
30 $\Delta A_{590}/g$) to the level in Plaisant 1 malt (1.617 $\Delta A_{590}/g$).

All analyses described above were carried out

at least in duplicate and the mean values are presented. The experimental error (E.E.) was calculated from the difference (in %) between the individual and the mean values.

5

SDS-Polyacrylamide gel electrophoresis and iso-electrofocusing

The molecular weight of the purified inhibitor was determined by SDS-polyacrylamide gel electrophoresis (SDS-page) on 20% polyacrylamide gels under reducing (β -mercaptoethanol, 1%) or non-reducing conditions with the PhastSystem unit (Pharmacia, Uppsala, Sweden), according to the method of Laemmli, U.K. (Nature, 1970, 227, 680-685). The gels were silver stained according to the instructions of the manufacturer (Pharmacia, Development Technique file N° 210). Low molecular weight markers were α -lactalbumin (14,400 Da); trypsin inhibitor (20,100 Da); carbonic anhydrase (30,000 Da); ovalbumin (43,000 Da); albumin (67,000 Da); phosphorylase b (94,000 Da). The iso-electric point of the inhibitor was determined with the PhastSystem unit using polyacrylamide gels containing ampholytes (pH 3-9) and with appropriate standards (Pharmacia calibration kits, pI 3.5 - 9.3). The proteins were silver stained (cfr. supra).

25

N-terminal amino acid sequencing of proteins.

The sequences of the N-terminal amino acids were determined with an Applied Biosystems model 477 A gas-phase sequencer, connected on line with an 120 A PTH analyser (Perkin Elmer, Belgium).

30

Evidence for the presence of endoxylanase inhibitors in wheat

Barley malt and wheat Xyl levels and arabinoxylan hydrolysing activities are listed in Table I. Xyl levels in the 100% malt worts (Table II) varied from 0.41 to 0.62% (all analytical data expressed as percentage of dry matter). The Xyl levels in the worts with 40% wheat varied from 0.35 to 0.61% (Table III).

In the worts with 40% wheat, the inventors used 60% barley malt. Comparison of the increase in Xyl during brewing using 60% malt with 60% of the Xyl increase using 100% barley malt showed a reduction of 12 to 58% (Tables II and III). This suggested that the endoxylanases from barley malt were inhibited in the presence of wheat or that the wheat arabinoxylans are a less suitable substrate for malt endoxylanases. Malting breaks down barley cell walls and renders them more accessible for enzymes (Selvig, A., & Aarnes, H., *Journal of the Institute of Brewing*, 1986, 92,185).

20

Free Xyl levels in wort

The levels of free Xyl in 100% malt worts varied from 0.046 to 0.076% and in the worts with 40% wheat from 0.025 to 0.040% (Table IV). The difference between the levels of the released free Xyl was 0.032 to 0.044% for the 100% malt worts and 0.015 to 0.020% for the worts with 40% wheat. The reduction in free Xyl release compared with 60% of the free Xyl release with the 100% barley malt wort varied from 1 to 32% (Tables II and IV). The use of the *Bacillus subtilis* endoxylanase did not result in an increase of free Xyl. The free Ara levels did also not increase. The endoxylanase, therefore, had no side β -D-

xylosidase and α -L-arabinofuranosidase activities.

The reduction of the endoxylanase induced increase in Xyl or arabinoxylan solubilization as a result of the use of wheat in conjunction with barley malt was more obvious than the reduction of the release of free Xyl. For this reason one focused on the inhibition of the barley malt endoxylanases by a wheat component.

Malt xylanolytic system inhibition by wheat extracts

10 In Figure 1 the reduction of the endoxylanase activity of BMWM1, when WWM instead of buffer A was added, is given. The reduction of endoxylanase activity varied from 26 to 58 %. The reduction of endoxylanase activity of barley malt wholemeal extracts (BMWM1) was obtained by
15 addition of wheat wholemeal extracts (WWM) instead of buffer. The figure 1 represents the results obtained with unboiled (\square) and boiled extracts (\square). (a) Clarine malt + Soissons wheat, (b) Clarine malt + Skirlou wheat, (c) Plaisant 1 malt + Soissons wheat, (d) Plaisant 1 malt +
20 Skirlou wheat, (e) Plaisant 2 malt + Soissons wheat, (f) Plaisant 2 malt + Skirlou wheat.

A higher reduction was observed in case of cv. Skirlou than with cv. Soissons. This was in line with the higher reduction of Xyl increase during brewing with
25 cv. Skirlou than with cv. Soissons (see Table III). The higher water extractable Xyl content of cv. Skirlou than for cv. Soissons implied that the lower susceptibility of the wheat substrate during brewing may cause the reduced solubilization. When boiling WWM, almost all of the
30 inhibition disappeared. The inhibitor seemed to be thermolabile and the inventors concluded that it therefore may be of proteic nature. However, it was considered

unlikely that one dealt with a protease as the protease activity from the malt is many times higher than the protease activity of wheat. The major part of the reduced activity was apparently not caused by the wheat
5 arabinoxylans because the thermal treatment did not change the arabinoxylan concentration of the wheat extract. Whether the wheat inhibitor was active against the endogenous barley malt endoxylanases or exogenous endoxylanases was unclear.

10

Brewing with Bacillus subtilis xylanase: a solution for poor arabinoxylan solubilisation in barley malt - wheat wholemeal brewing

The Bacillus subtilis endoxylanase, one of
15 the microbial enzymes that was relatively little inhibited under the experimental conditions of Figure 2, increased the Xyl levels present in wort. In comparison with the same worts made without endoxylanase addition, 94% more Xyl solubilization using cv. Soissons as wheat adjunct and 179%
20 more Xyl solubilization using cv. Skirlou as wheat adjunct was obtained. The Bacillus subtilis endoxylanase apparently solubilizes more arabinoxylan from the wheat variety Skirlou than from the wheat variety Soissons (Table IV).

25 Purification of xylanase inhibitor from wheat flour

Soissons wheat flour (2.0 kg) was suspended in 10.0 L 0.1% (w/v) ascorbic acid. The suspension was mixed overnight at 7°C and centrifuged (7°C, 10,000 g, 30 min). To the supernatant 2.0 g/l CaCl₂ was added and the pH
30 was raised to 9.0 by addition of 2.0 M NaOH. The extract was left overnight at 7°C and centrifuged (7°C, 10,000 g, 30 min). The pH of the extract was adjusted to 5.0 with 2.0

M HCl and the extract was pumped over a cation exchanger (SP Sepharose Fast Flow, 50 x 50 mm, Pharmacia). The column was equilibrated with buffer C (200 mL) and a protein fraction was eluted with 200 mL 0.5 M NaCl. This eluate was
5 diluted 5 times, the pH adjusted to 5.0 as above and cations were exchanged (SP Sepharose Fast Flow, 26 x 100 mm, Pharmacia). The column was equilibrated with buffer C (200 mL) and after a linear salt gradient from 0 to 0.5 M NaCl (800 mL), fractions of 10 mL were, after desalting (PD
10 10 column, Pharmacia), assayed for endoxylanase inhibition using the cited xylazyme method with eluate instead of cereal extract and appropriately diluted Xylanase M4 from *Aspergillus niger*. The fractions with inhibition activity were collected, dialyzed against deionised water (7°C,
15 overnight) and lyophilised. The lyophilised material was dissolved in buffer D (6.0 mL) and separated on a Sephacryl S100 column (26 x 670 mm, Pharmacia) eluted with the same buffer. Fractions of 2.5 mL were collected and assayed for inhibitor activity. The active fractions were collected,
20 dialyzed as above and lyophilised. The lyophilised material was dissolved in buffer E (6 mL) and cation exchanged (Mono S HR 5/5, Pharmacia) with the same buffer. Fractions eluted in a salt gradient (0 to 0.5 M NaCl) were collected and assayed for xylanase inhibition as above. In this way,
25 we obtained a fraction (1 mL) of the inhibitor which migrated as a single protein band on SDS-PAGE. It had an apparent molecular weight of ca. 40 - 43 kDa. Following reduction with β -mercaptoethanol, two additional SDS-PAGE bands of molecular weights of typically 30 and 10 kDa are
30 detected.

N-terminal amino acid sequencing of endoxylanase inhibitor

The β -mercaptoethanol reduced inhibitor protein and/or glycoprotein fractions were subjected to SDS-page, blotted and N-terminal amino acid sequenced.

5 The N-terminal amino acid sequence of the ca. 40 - 43 kDa band (SEQ ID No.01) was: Lys-Gly-Leu-Pro-Val-Leu-Ala-Pro-Val-Thr-Lys-Xaa-Thr-Ala wherein Xaa being preferably Asp. This sequence has not been reported before. The above cited ca. 30 kDa band also has the above
10 described typical N-terminal amino acid SEQ ID No.1, while the N-terminal amino acid sequence of the ca. 10 kDa band is typically as follows: SEQ ID No. 2: Xaa-Ala-Pro-Val-Ala-Lys-Met-Val-Leu-Pro-Val-Ala-Met-Lys-Glu-Xaa-Val, wherein the first Xaa being preferably Ser, Phe, or Gly, and
15 wherein the second Xaa is unidentified. This sequence has not been described before.

Inhibition of different microbial endoxylanases by endoxylanase inhibitors from wheat and other cereals

20 In Figure 2 the inhibition of different microbial xylanases in the presence of WF, RF, and BWM is shown. The reduction of the xylanase activity (%) when a cereal extract was added instead of the same extract boiled for 30 min is given. Under the experimental conditions, the
25 highest reduction was found for the mixture of three xylanases from *Trichoderma reesei* (82 to 94%) the lowest for the xylanases from *Bacillus subtilis* (24 to 39%).

 The reduction of microbial endoxylanase activity was obtained by addition of cereal extracts (WF, RF, and BWM) instead of boiled cereal extracts. The figure
30 2 represents the results obtained with wheat flour (■),

rye flour (□) and barley whole meal (■). Microbial xylanases: (a) Mixture of three xylanases from *Trichoderma reesei*, (b) Xylanase M4 from *Aspergillus niger*, (c) Xylanase from *Bacillus subtilis*, (d) Mixture of three
5 xylanases from *Bacillus subtilis*, (e) Xylanase from *Aspergillus niger*, (f) Mixture of five xylanases from *Aspergillus niger*, (g) Mixture of five xylanases from *Aspergillus niger*. Under the experimental conditions, WG reduced the activity of xylanase M4 from *Aspergillus niger*
10 with ca. 80 %.

Inhibition of barley malt β -glucanase by inhibitors from wheat and other cereals

In Figure 3 the inhibition of malt β -
15 glucanase in the presence of WF, RF, and BWM is shown. The reduction of the β -glucanase activity (%) when a cereal extract was added instead of the same extract boiled for 30 min is given. The reduction varied from 7 to 12%.

The reduction of β -glucanase activity of
20 barley malt extracts (BMWM2) was obtained by addition of cereal extracts (WF, RF, BWM) instead of boiled cereal extracts. The figure 3 represents the results obtained with wheat flour (a), rye flour (b) and barley whole meal (c).

TABLE I. Water -extractible and Free Xylose Contents (% of Dry Matter) and Arabinoxylan Degrading Enzyme Activities of 3 Barley Malts and 2 Wheats*.

| | Water-extractable Xyl | Free Xyl | Endoxylanase ($\Delta A_{590}/g$) | β -D-Xylosidase (U/g) |
|-------------|-----------------------|----------|--|--------------------------------|
| Barley Malt | | | | |
| Clarine | 0.29 | 0.014 | 0.750 | 0.286 |
| Plaisant 1 | 0.41 | 0.031 | 1.617 | 0.331 |
| Plaisant 2 | 0.40 | 0.013 | 0.607 | 0.299 |
| Wheat | | | | |
| Soissons | 0.27 | 0.005 | 0.115 | 0.054 |
| Skirlou | 0.52 | 0.003 | 0.205 | 0.053 |
| | E.E.<7% | E.E.<4% | E.E. <9% | E.E. <6% |

*Xyl = xylose; Water-extractable Xyl = total - free xylose water-extract; E.E. = experimental error.

TABLE II Xylose and Free Xylose Levels (% of Dry Matter) in Wort and Levels of Increase in Xylose and Free Xylose (% of Dry Matter) during Brewing with 100% Barley Malt*.

| Barley Malt | Xyl Wort | Xyl Increase | Free Xyl Wort | Free Xyl Increase |
|-------------|----------|--------------|---------------|-------------------|
| Clarine | 0.41 | 0.15 | 0.046 | 0.032 |
| Plaisant 1 | 0.62 | 0.26 | 0.075 | 0.044 |
| Plaisant 2 | 0.51 | 0.09 | 0.049 | 0.035 |

23

E.E.<6%

E.E.<7%

*Xyl = xylose; Xyl Wort = total - free xylose wort; Xyl Increase = total xylose wort - total xylose water-extract barley malt; Ara = arabinose; E.E. = experimental error.

TABLE III

Xylose Levels (% of Dry Matter) in Grains and Corresponding Worts Prepared with 60% Barley Malt and 40% Wheat. Increase in Xylose Levels (% of Dry Matter) during Brewing and Effect of Addition of *Bacillus subtilis* Endoxylanase. Difference (%) with 60% of the Increase of Xylose Levels in Case of a 100% Malt Wort*.

| Barley Malt (60%) + Wheat (40%) | Xyl Grains | Xyl Wort | Xyl Increase | Difference 100% Malt Wort |
|------------------------------------|------------|----------|--------------|------------------------------|
| Clarine + Soissons | 0.29 | 0.35 | 0.08 | -12 |
| Clarine + Skirlou | 0.41 | 0.46 | 0.07 | -27 |
| Plaisant 1 + Soissons | 0.35 | 0.44 | 0.10 | -36 |
| Plaisant 1 + Skirlou | 0.48 | 0.53 | 0.06 | -58 |
| Plaisant 2 + Soissons | 0.35 | 0.39 | 0.06 | -28 |
| Plaisant 2 + Skirlou | 0.48 | 0.51 | 0.05 | -40 |
| Clarine + Soissons + BSX | 0.31 | 0.45 | 0.16 | 93 |
| Clarine + Skirlou + BSX | 0.44 | 0.61 | 0.19 | 130 |
| E.E.<6% | | | | E.E.<7% |

$\text{*Xyl} = \text{xylose}; \text{Xyl Grains} = 0.6 \times (\text{total} - \text{free xylose water-extract barley malt}) + 0.4 \times$
 $(\text{total} - \text{free xylose water-extract wheat}); \text{Xyl wort} = \text{total} - \text{free xylose wort}; \text{Xyl}$
 $\text{Increase} = \text{total xylose wort} - [0.6 \times (\text{total xylose water-extract barley malt}) + 0.4 \times$
 $(\text{total xylose water-extract wheat})]; \text{Difference } 100\% \text{ Malt Wort} = [100 \times (\text{increase xylose}$
 $\text{wort from } 60\% \text{ malt and } 40\% \text{ wheat}) / (\text{xylose increase wort from } 100\% \text{ malt})] - 100; \text{BSX} =$
 $\text{Bacillus subtilis endoxylanase}; \text{E.E.} = \text{experimental error}.$

TABLE IV

Release of Free Xylose during Brewing with 60% Barley Malt and 40% Wheat. Effect of Addition of *Bacillus subtilis* Endoxylanase. Difference (%) with 60% of the Release of Xylose in Case of a 100% Malt Wort*.

| Free Xyl | | | |
|--------------------------|---------|---------|-------------------------------|
| Barley Malt (60%) + | Grains | Wort | Release |
| Wheat (40%) | | | Difference 100 % Malt Wort |
| Clarine + Soissons | 0.010 | 0.025 | 0.015 |
| Clarine + Skirlou | 0.010 | 0.029 | 0.019 |
| Plaisant 1 + Soissons | 0.021 | 0.039 | 0.018 |
| Plaisant 1 + Skirlou | 0.020 | 0.040 | 0.020 |
| Plaisant 2 + Soissons | 0.010 | 0.029 | 0.019 |
| Plaisant 2 + Skirlou | 0.009 | 0.029 | 0.020 |
| Clarine + Soissons + BSX | 0.010 | 0.026 | 0.015 |
| Clarine + Skirlou + BSX | 0.010 | 0.029 | 0.019 |
| | | | |
| | E.E.<8% | E.E.<7% | |

*Xyl = xylose; Difference 100% Malt Wort = $[100 \times (\text{xylose release wort from 60\% malt and 40\% wheat}) / (\text{xylose release wort from 100\% malt})] - 100$; BSX = *Bacillus subtilis* endoxylanase; E.E. = experimental error.

Claims

1. A proteinic or glycoprotein inhibitor of cellulolytic, xylanolytic and/or β -glucanolytic enzymes.

2. Inhibitor as in claim 1, characterised in that said inhibitor inhibits cellulase, endoxylanase, β -glucanase, β -xylosidase, α -L-arabinofuranosidase and/or other cellulose, xylan, arabinoxylan or β -glucan degrading enzymes.

3. Inhibitor as in claim 1 or 2, characterised in that it is obtainable from plant material or fractions thereof.

4. Inhibitor as in claim 3, characterised in that said plant material is selected from the group consisting of cereals, cereal grains or cereal flours from wheat, durum wheat, rye, triticale, barley, sorghum, oats, maize or rice.

5. Inhibitor as in claim 1 or 2, characterised in that it is obtainable from micro-organisms or fractions thereof.

6. Inhibitor as in any of the claims 1-5, characterised in that it is a xylanase inhibitor.

7. Inhibitor as in claim 6, characterised in that it is a water-soluble species.

8. Inhibitor as in claim 7 having a marker whose amino acid sequence has more than 70% homology with SEQ ID No. 1 and/or SEQ ID No.2.

9. Inhibitor as in claim 8, characterised in that the marker is the N-terminal amino acid sequence of

the protein or glycoprotein.

10. Inhibitor as in claim 8 or 9 having a marker for which amino acid sequence has more than 85% homology with SEQ ID No. 1 and/or SEQ ID No.2.

11. Inhibitor as in claim 10, characterised in that the marker is the N-terminal amino acid sequence of the protein or glycoprotein.

12. Inhibitor as in claim 7, having a marker whose amino acid sequence is identical to SEQ ID No. 1 and/or SEQ ID No.2.

13. Inhibitor as in claim 12, characterised in that the marker is the N-terminal amino acid sequence of the protein or glycoprotein.

14. Inhibitor as in any of the claims 7 to 13, characterised in that said protein or glycoprotein is selected from the group comprising proteins or glycoproteins having a molecular weight typically between 40 kDa and 43 kDa, proteins or glycoproteins having a molecular weight typically 30 kDa and proteins or glycoproteins having a molecular weight of typically 10 kDa.

15. Inhibitor as in any of the claims 7 to 14, characterised in that said protein or glycoprotein typically a molecular weight between 40 kDa and 43 kDa and has a pI of greater than about 7.

16. Method for obtaining the inhibitor as in any of the claims 1 to 15 from possibly genetically modified micro-organisms, plants or plant materials, wherein said micro-organisms, plants or plant materials are subjected to one or more extraction and/or fractionation

steps.

17. Method for obtaining the inhibitor according to any of the claims 1 to 15, wherein micro-organisms, plants or plant materials are genetically modified by the introduction of a genetic material encoding said inhibitor into the micro-organisms, plants or plant materials.

18. Process for transforming micro-organisms, plants or plant materials, wherein the activity of the inhibitor according to any of the claims 1 to 15 is reduced.

19. Process according to claim 18, characterised in that the reduced activity of the inhibitor according to the invention is obtained by reduction of its expression.

20. Process according to claim 18 or 19, characterised in that the activity of the inhibitor is reduced by blocking the inhibitor function.

21. Process for transforming micro-organisms, plants or plant materials, wherein the activity of the inhibitor according to any of the claims 1 to 15 is increased.

22. Process according to claim 21, characterised in that increased activity of the inhibitor according to the invention is obtained by an increase of its expression.

23. Process according to claim 21 or 22, characterised in that the activity of the inhibitor is increased by activating the inhibitor function.

24. Micro-organisms, plants or plant

materials obtained by the method according to any of the preceding claims 17 to 23.

25. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants and/or the plant materials according to claim 24 for improving the malting of cereals such as barley, sorghum and wheat and/or the production of beer.

26. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant materials according to claim 24 for improving the production and/or quality of baked or extruded cereal products chosen among the group consisting of straight dough, sponge dough, Chorleywood bread, breakfast cereals, biscuits, pasta and noodles.

27. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant materials according to claim 24 for improving animal feedstuff efficiency.

28. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant materials according to claim 24 for improving the production of starch derived syrups, sorbitol, xylose and/or xylitol.

29. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant

31

materials according to claim 24 for wheat gluten-starch separation and production.

30. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant materials according to claim 24 for improving maize processing.

31. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant materials according to claim 24 for improving plant disease resistance.

32. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant materials according to claim 24 for improving nutraceutical and/or pharmaceutical applications.

33. Use of the inhibitor according to any of the preceding claims 1 to 15 or obtained by the method of claim 16, the micro-organisms, the plants or the plant materials according to claim 24 for improving paper and pulp technologies.




Fig. 1




Fig. 2




Fig. 3

2/3
SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: K.U. LEUVEN RESEARCH & DEVELOPMENT
(B) STREET: Groot Begijnhof 59
(C) CITY: Leuven
(E) COUNTRY: BELGIUM
(F) POSTAL CODE (ZIP): B-3000

(A) NAME: DEBYSER Winok
(B) STREET: Willem Coosemansstraat 23
(C) CITY: Kessel-Lo
(E) COUNTRY: BELGIUM
(F) POSTAL CODE (ZIP): B-3010

(A) NAME: DELCOUR Jan
(B) STREET: Kastanjelaan, 14
(C) CITY: Leuven
(E) COUNTRY: BELGIUM
(F) POSTAL CODE (ZIP): B-3001

(ii) TITLE OF INVENTION: INHIBITORS OF CELLULOLYTIC, XYLANOLYTIC AND
BETA-GLUCANOLYTIC ENZYMES

(iii) NUMBER OF SEQUENCES: 2

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 12
(D) OTHER INFORMATION: /product= "preferably Asp"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Lys Gly Leu Pro Val Leu Ala Pro Val Thr Lys Xaa Thr Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

3/3

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION:1
- (D) OTHER INFORMATION:/product= "Ser, Phe, or Gly"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION:16
- (D) OTHER INFORMATION:/product= "Unidentified"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Xaa | Ala | Pro | Val | Ala | Lys | Met | Val | Leu | Pro | Val | Ala | Met | Lys | Glu | Xaa |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |

Val

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 98/02590

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/24 A23L1/185 A23L1/10 C07K14/415

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A23L C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| Y | ZISER, LOTHAR ET AL: "Syntheses and testing of substrates and mechanism -based inactivators for xylanases" CARBOHYDR. RES. (1995), 274, 137-53 CODEN: CRBRAT; ISSN: 0008-6215, XP002078666 whole document, esp. Figs. 1 and 2, Abstract | 1-33 |
| Y | KESKAR, SULABHA S. ET AL: "Characterization and sequencing of an active-site cysteine-containing peptide from the xylanase of a thermotolerant Streptomyces" BIOCHEM. J. (1992), 281(3), 601-5 CODEN: BIJOAK; ISSN: 0306-3275, XP002078667 Seite 603, Figs. 1-3 | 1-33 |

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "S" document member of the same patent family

Date of the actual completion of the international search

28 September 1998

Date of mailing of the international search report

13/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kronester-Frei, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/02590

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|--|-----------------------|
| Y | GOMES, D. J. ET AL: "Factors influencing the induction of endoxylanase by <i>Thermoascus aurantiacus</i> " J. BIOTECHNOL. (1994), 33(1), 87-94 CODEN: JBITD4; ISSN: 0168-1656, XP002078668 Abstract, Tables 1 and 2 --- | 1-33 |
| Y | PAUL, JAISHREE ET AL: "Influence of sugars on endoglucanase and beta.-xylanase activities of a <i>Bacillus</i> strain" BIOTECHNOL. LETT. (1990), 12(1), 61-4 CODEN: BILED3; ISSN: 0141-5492, XP002078669 Abstract and Fig. 2 --- | 1-33 |
| P,X | DEBYSER, WINOK ET AL: "Arabinoxylan solubilization and inhibition of the barley malt xylanolytic system by wheat during mashing with wheat wholemeal adjunct: evidence for a new class of enzyme inhibitors in wheat" J. AM. SOC. BREW. CHEM. (1997), 55(4), 153-156 CODEN: JSBCD3; ISSN: 0361-0470, XP002078670 Last sentence of Abstract and Conclusion --- | 1-6 |
| P,X | SPURWAY, TRACEY D. ET AL: "Calcium protects a mesophilic xylanase from proteinase inactivation and thermal unfolding" J. BIOL. CHEM. (1997), 272(28), 17523-17530 CODEN: JBCHA3; ISSN: 0021-9258, XP002078671 whole document, especially Fig. 8 and Abstract --- | 1-6 |
| P,Y | WO 98 05788 A (SOUPPE JEROME ;MOGEN INT (NL); BEUDEKER ROBERT FRANCISCUS (NL); GI) 12 February 1998 claims, examples --- | 1-33 |
| A | WO 96 29416 A (MADRID SUSAN MAMPUSTA ;DANISCO (DK); BARUCH ANITA (DK); RASMUSSEN) 26 September 1996 claims, examples ----- | 1-33 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/02590

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9805788 A | 12-02-1998 | AU 4116197 A | 25-02-1998 |
| WO 9629416 A | 26-09-1996 | AU 5104396 A | 08-10-1996 |
| | | BR 9607535 A | 06-01-1998 |
| | | CA 2214591 A | 26-09-1996 |